arXiv:0910.2766v2 [math.CO] 25 Jul 2011

PACKING 3-VERTEX PATHS
IN CUBIC 3-CONNECTED GRAPHS

Alexander Kelmans

University of Puerto Rico, San Juan, Puerto Rico
Rutgers University, New Brunswick, New Jersey

The paper was accepted for publication
in Discrete Mathematics in August 14, 2009

Abstract

A subgraph (a spanning subgraph) of a graph G whose all components are 3-vertex paths is called an Λ-packing (respectively, an Λ-factor of G. Let $v(G)$ and $\lambda(G)$ be the number of vertices and the maximum number of disjoint 3-vertex paths in G, respectively. We discuss the following old

Problem (A. Kelmans, 1984). *Is the following claim true?*
(P) *If* G *is a cubic 3-connected graph, then* G *has a* Λ*-packing that avoids at most two vertices of* G*, i.e.* $\lambda(G) = \lfloor v(G)/3 \rfloor$.

We show, in particular, that claim (P) is equivalent to some seemingly stronger claims (see **3.1**). For example, if $v(G) = 0 \bmod 3$, then the following claims are equivalent: G has a Λ-factor, for every $e \in E(G)$ there exists a Λ-factor of G avoiding (containing) e, $G - X$ has a Λ-factor for every $X \subseteq E(G)$ such that $|X| = 2$, and $G - L$ has a Λ-factor for every 3-vertex path L in G. We also show that certain claims in theorem **3.1** are best possible. We give a construction providing infinitely many cyclically 6-connected graphs G with two disjoint 3-vertex paths L and L' such that $v(G) = 0 \bmod 3$ and $G - L - L'$ has no Λ-factor. It turns out that if claim (P) is true, then Reed's dominating graph conjecture is true for cubic 3-connected graphs.

Keywords: cubic 3-connected graph, 3-vertex path packing, 3-vertex path factor, domination.

1 Introduction

We consider undirected graphs with no loops and no parallel edges. All notions and facts on graphs, that are used but not described here, can be found in [1, 2, 19].

Given graphs G and H, an *H-packing* of G is a subgraph of G whose all components are isomorphic to H. An *H-packing* P of G is called an *H-factor* if $V(P) = V(G)$. The *H-packing problem*, i.e., the problem of finding in G an H-packing, having the maximum number of vertices, turns out to be NP-hard for every connected graph H with at least three vertices [3]. Let Λ denote a 3-vertex path. In particular, the Λ-packing problem is NP-hard. Moreover, the problem of packing k-vertex paths, $k \ge 3$, is NP-hard for cubic graphs [6]. Also, the 3-vertex path packing problem remains NP-hard even for cubic bipartite planar graphs [16].

Although the Λ-packing problem is NP-hard, i.e., possibly intractable in general, this problem turns out to be tractable for some natural classes of graphs. It would be also interesting to find polynomial algorithms that would provide a good approximation for the problem. Below (see **1.1 - 1.2**) there are some examples of such results. In each case the corresponding packing problem is polynomially solvable.

Let $v(G)$ and $\lambda(G)$ denote the number of vertices and the maximum number of disjoint 3-vertex paths in G, respectively. Obviously, $\lambda(G) \le \lfloor v(G)/3 \rfloor$.

A graph is called *claw-free* if it contains no induced subgraph isomorphic to $K_{1,3}$ (which is called a *claw*). A *block* of a graph G is either an isolated vertex or a maximal connected subgraph H of G such that $H - v$ is connected for every vertex v of H. A block of a connected graph is called an *end-block* if it has at most one vertex in common with any other block of the graph. Let $eb(G)$ denote the number of end-blocks of G.

1.1 [12] *Suppose that G is a connected claw-free graph and $eb(G) \ge 2$. Then $\lambda(G) \ge \lfloor (v(G) - eb(G) + 2)/3 \rfloor$, and this lower bound is sharp.*

1.2 [12] *Suppose that G is a connected and claw-free graph having at most two end-blocks (in particular, a 2-connected and claw-free graph). Then $\lambda(G) = \lfloor v(G)/3 \rfloor$.*

Obviously, the claim in **1.2** on claw-free graphs with exactly two end-blocks follows from **1.1**.

In [5, 13] we answered the following natural question:

How many disjoint 3-vertex paths must a cubic graph have?

1.3 *If G is a cubic graph, then $\lambda(G) \ge \lceil v(G)/4 \rceil$. Moreover, there is a polynomial time algorithm for finding a Λ-packing having at least $\lceil v(G)/4 \rceil$ components.*

Obviously, if every component of G is K_4, then $\lambda(G) = v(G)/4$. Therefore the bound in **1.3** is sharp. There are some other connected cubic graphs for which the bound in **1.3** is attained.

Let $\mathcal{G}_2^3$ denote the set of graphs with each vertex of degree at least 2 and at most 3. In [5] we answered (among other results) the following question:

How many disjoint 3-vertex paths must an n-vertex graph from $\mathcal{G}_2^3$ have?

1.4 *Suppose that $G \in \mathcal{G}_2^3$ and G has no 5-vertex components. Then $\lambda(G) \ge \lceil v(G)/4 \rceil$.*

Obviously, **1.3** follows from **1.4** because if G is a cubic graph, then $G \in \mathcal{G}_2^3$ and G has no 5-vertex components.

In [5] we also gave a construction that allowed to prove the following:

1.5 *There are infinitely many connected graphs for which the bound in* **1.4** *is attained. Moreover, there are infinitely many subdivisions of cubic 3-connected graphs for which the bound in* **1.4** *is attained.*

The next interesting question is:

How many disjoint 3-vertex paths must a cubic connected graph have?

The following bound was recently found in [14].

1.6 *If G is a cubic connected graph and $v(G) \ge 17$, then $\lambda(G) \ge \frac{39}{152}v(G)$.*

The next natural question is:

1.7 Problem. *How many disjoint 3-vertex paths must a cubic 2-connected graph have?*

1.8 [15] *If G is a cubic connected graph and $v(G) \ge 9$, then $\lambda(G) \ge \frac{3}{11}v(G)$.*

On the other hand, it is also natural to consider the following

1.9 Problem. *Are there 2-connected cubic graphs G such that $\lambda(G) < \lfloor v(G)/3 \rfloor$?*

In [7] we gave a construction that provided infinitely many 2-connected, cubic, bipartite, and planar graphs such that $\lambda(G) < \lfloor v(G)/3 \rfloor$.

The main goal of this paper (see also [8]) is to discuss the following old open problem which is similar to Problem **1.9**.

1.10 Problem. (A. Kelmans, 1984). *Is the following claim true?*

(P) *if G is a 3-connected and cubic graph, then $\lambda(G) = \lfloor v(G)/3 \rfloor$.*

Originally we have we've become interested in this problem because it is related with non-trivial relaxations of some graph Hamiltonicity problems we've been studying (see [4]).

Problem **1.10** is also related with the minimum domination problem in a graph. In [18] B. Reed conjectured that if G is a connected cubic graph, then $\gamma(G) \le \lceil v(G)/3 \rceil$, where $\gamma(G)$ is the dominating number of G (i.e., the size of a minimum vertex subset X in G such that every vertex in $G - X$ is adjacent to a vertex in X). It turns out that Reed's conjecture is not true for connected and even for 2-connected cubic graphs [10,17]. If claim (P) in **1.10** is true, then from **3.1** it follows, in particular, that Reed's conjecture is true for 3-connected cubic graphs. In [9] we proved that if G is a cubic Hamiltonian graph with $v(G) = 1 \bmod 3$, then $\gamma(G) \le \lfloor v(G)/3 \rfloor$.

In this paper we show, in particular, that claim (P) in **1.10** is equivalent to some seemingly stronger claims (see **3.1**).

In Section 2 we give some notation, constructions, and simple observations. In Section 3 we formulate and prove our main theorem **3.1** concerning various claims that are



Figure 1: $Aa\sigma bB$

equivalent to claim (P) in **1.10**. We actually give different proofs of **3.1**. Thus, if there is a counterexample C to one of the above claims, then these different proofs provide different constructions of counterexamples to the other claims in **3.1**. Additionally, different proofs provide better understanding of relations between various Λ-packing properties considered in **3.1**. In Section 4 we describe some properties of Λ-factors with respect to 3-edge cut-matchings and triangles in cubic 3-connected graphs. In Section 5 we give two Λ-factor homomorphism theorems for cubic graphs. Finally, in Section 6 we give a construction of cubic cyclically 6-connected graphs showing that certain claims in **3.1** are best possible.

We can also prove that many equivalences in **3.1** are also true in the class of bipartite 3-connected cubic graphs and that if claim (P) in **1.10** is true for bipartite graphs, then Reed's conjecture is also true for bipartite 3-connected cubic graphs.

2 Notation, constructions, and simple observations

We consider undirected graphs with no loops and no parallel edges unless stated otherwise. As usual, $V(G)$ and $E(G)$ denote the set of vertices and edges of G, respectively, and $v(G) = |V(G)|$. If X is a vertex subset or a subgraph of G, then let $D(X, G)$ (or simply $D(X)$) denote the set of edges in G having exactly one end-vertex in X, and let $d(X, G) = |D(X, G)|$. If $x \in V(G)$, then $D(x, G)$ is the set of edges in G incident to x, $d(x, G) = |D(x, G)|$, $N(x, G) = N(x)$ is the set of vertices in G adjacent to x, and $\Delta(G) = \max\{d(x, G) : x \in V(G)\}$. If $e = xy \in E(G)$, then let $End(e) = \{x, y\}$. Let $Cmp(G)$ denote the set of components of G and $cmp(G) = |Cmp(G)|$.

We call an edge cut K in a cubic graph a *blockade*, if K is matching. A *k-blockade* is a blockade with k edges. Every connected cubic graph non-isomorphic to K_4 or $K_{3,3}$ has a blockade. For every such graph G, let $c(G)$ denote the size of a minimum blockade in G; the parameter $c(G)$ is called the *cyclic connectivity of* G and G is called *cyclic k-connected*, if $c(G) \ge k$.

Let A and B be disjoint graphs, $a \in V(A)$, $b \in V(B)$, and $\sigma : N(a, A) \to N(b, B)$ be a bijection. Let $Aa\sigma bB$ denote the graph $(A - a) \cup (B - b) \cup \{x\sigma(x) : x \in N(a, A)\}$. We usually assume that $N(a, A) = \{a_1, a_2, a_3\}$, $N(b, B) = \{b_1, b_2, b_3\}$, and $\sigma(a_i) = b_i$ for $i \in \{1, 2, 3\}$ (see Fig. 1). We also say that *$Aa\sigma bB$ is obtained from B by replacing* *vertex b by $(A - a)$ according to σ.*

Let B be a cubic graph and $X \subseteq V(B)$. Let A^v, where $v \in X$, be a graph, a^v be a vertex of degree three in A^v, and $A_v = A^v - a^v$. By using the above operation, we can build a graph $G = B\{(A^v, a^v) : v \in X\}$ by replacing each vertex v of B in X by A_v assuming that all A^v's are disjoint. We call $G = B\{(A^v, a^v) : v \in V(B)\}$ a *graph-composition*, B – the *frame* and each (A^v, a^v) – a *brick* of this graph-composition. Let $D^v = D(A_v, G)$. If all (A^v, a^v)'s are isomorphic to some (Z, z), then instead of $B\{(A^v, a^v) : v \in V(B)\}$ we simply write $B\{Z, z\}$. For each $u \in V(B) \setminus X$, let A^u be the graph having exactly two vertices u, a^u and exactly three parallel edges connecting u and a^u. Then $G = B\{(A^v, a^v) : v \in X\} = B\{(A^v, a^v) : v \in V(B)\}$. If, in particular, $X = V(B)$ and each A^v is a copy of K_4, then G is obtained from B by replacing each vertex by a triangle.

Let $E' = E'(G) = E(G) \setminus \cup\{E(A_v) : v \in V(B)\}$. Obviously, there is a unique bijection $\alpha : E(B) \to E'(G)$ such that if $uv \in E(B)$, then $\alpha(uv)$ is an edge in G having one end-vertex in A_u and the other in A_v.

Let P be a Λ-packing in G. For $uv \in E(B)$, $u \neq v$, we write $u \neg^P v$ or simply, $u \neg v$, if P has a 3-vertex path L such that $\alpha(uv) \in E(L)$ and $|V(A_u) \cap V(L)| = 1$. Let P^v be the union of components of P that meet (i.e., have an edge in) D^v.

Obviously:

2.1 *Let k be an integer and $k \leq 3$. If A and B above are k-connected, cubic, bipartite, and planar graphs, then $Aa\sigma bB$ is also a k-connected, cubic, bipartite, and planar graph, respectively.*

From **2.1** we have:

2.2 *Let k be an integer and $k \leq 3$. If B and each A^v is a k-connected, cubic, bipartite, and planar graphs, then $B\{(A^v, a^v) : v \in V(B)\}$ is also a k-connected, cubic, bipartite, and planar graph, respectively.*

Let A^1, A^2, A^3 be three disjoint graphs, $a^i \in V(A^i)$, and $N(a^i, A^i) = \{a^i_1, a^i_2, a^i_3\}$, where $i \in \{1, 2, 3\}$. Let $F = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ denote the graph obtained from $(A^1 - a^1) \cup (A^2 - a^2) \cup (A^3 - a^3)$ by adding three new vertices z_1, z_2, z_3 and the set of nine new edges $\{z_j a^i_j : i, j \in \{1, 2, 3\}$ (see Fig. 2). In other words, if $B = K_{3,3}$ is the complete (X, Z)-bipartite graph with $X = \{x_1, x_2, x_3\}$ and $Z = \{z_1, z_2, z_3\}$, then F is obtained from B by replacing each vertex x_i in X by $A^i - a^i$ so that $D(A^i - a^i, F) = \{a^i_j z_j : j \in \{1, 2, 3\}\}$. Let $D^i = D(A^i - a^i, F)$. If P is a Λ-packing of F, then let $P^i = P^i(F)$ be the union of components of P meeting D^i and $E^i = E^i(P) = E(P) \cap D^i$, $i \in \{1, 2, 3\}$.

If each (A^i, a^i) is a copy of the same (A, a), then we write $Y(A, a)$ instead of $Y(A^1, a^1; A^2, a^2; A^3, a^3)$.

In particular, from **2.2** we have:



Figure 2: $Y(A^1, a^1; A^2, a^2; A^3, a^3)$



Figure 3:

2.3 *Let k be an integer and $k \leq 3$. If each A^i above is a k-connected, cubic, and bipartite graph, then $Y(A^1, a^1; A^2, a^2; A^3, a^3)$* (see Fig. 2) *is also a k-connected, cubic, and bipartite graph, respectively.*

We will also use the following simple observation.

2.4 *Let A and B be disjoint graphs, $a \in V(A)$, $N(a, A) = \{a_1, a_2, a_3\}$, $b \in V(B)$, $N(b, B) = \{b_1, b_2, b_3\}$, and $G = Aa\sigma bB$, where each $\sigma(a_i) = b_i$* (see Fig. 1). *Let P be a Λ-factor of G (and so $v(G) = 0 \bmod 3$) and P' be the Λ-packing of G consisting of the components (3-vertex paths) of P that meet $\{a_1b_1, a_2b_2, a_3b_3\}$.*

(*a*1) *Suppose that $v(A) = 0 \bmod 3$, and so $v(B) = 2 \bmod 3$. Then one of the following holds* (see Fig. 3):

(*a*1.1) *P' has exactly one component that has two vertices in $A - a$, that are adjacent (and, accordingly, exactly one vertex in $B - b$),*

(*a*1.2) *P' has exactly two components and each component has exactly one vertex in $A - a$ (and, accordingly, exactly two vertices in $B - b$, that are adjacent),*

(*a*1.3) *P' has exactly three components L_1, L_2, L_3 and one of them, say L_1, has exactly one vertex in $A - a$ and each of the other two L_2 and L_3 has exactly two vertices in $A - a$, that are adjacent (and, accordingly, L_1 has exactly two vertices in $B - b$, that are adjacent, and each of the other two L_2 and L_3 has exactly one vertex in $B - b$).*

(*a*2) *Suppose that $v(A) = 1 \bmod 3$, and so $v(B) = 1 \bmod 3$. Then one of the following holds* (see Fig. 4)*:*



Figure 4:

($a2.1$) $P' = \emptyset$,

($a2.2$) P' has exactly two components, say L_1 and L_2, such that one of the them, say L_1, has exactly one vertex in $A - a$ and exactly two vertices in $B - b$, that are adjacent, and the other component L_2 has exactly two vertices in $A - a$, that are adjacent, and exactly one vertex in $B - b$,

($a2.3$) P' has exactly three components L_1, L_2, L_3 and either each L_i has exactly one vertex in $A - a$ (and, accordingly, has exactly two vertices in $B - b$, that are adjacent) or each L_i has exactly two vertices in $A - a$, that are adjacent (and, accordingly, has exactly one vertex in $B - b$).

3 Λ-packings in cubic 3-connected graphs

The main goal of this section is to prove the following theorem showing that claim (P) in **1.10** is equivalent to various seemingly stronger claims.

3.1 *The following claims are equivalent for cubic 3-connected graphs G:*

(**z1**) $v(G) = 0 \bmod 6 \Rightarrow$ *G has a Λ-factor,*

(**z2**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every $e \in E(G)$ there is a Λ-factor of G avoiding e (i.e., $G - e$ has a Λ-factor),*

(**z3**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every $e \in E(G)$ there exists a Λ-factor of G containing e,*

(**z4**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exists at least one 3-vertex path L such that L is centered at x and $G - L$ has a Λ-factor,*

(**z5**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exist at least two 3-vertex paths L such that L is centered at x and $G - L$ has a Λ-factor,*

(**z6**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every $xy \in E(G)$ there exist edges $xx', yy' \in E(G)$ such that*

$G - xyy'$ and $G - x'xy$ have Λ-factors,

(**z7**) $v(G) = 0 \bmod 6 \Rightarrow$ *$G - X$ has a Λ-factor for every $X \subseteq E(G)$ such that $|X| = 2$,*

(**z8**) $v(G) = 0 \bmod 6 \Rightarrow$ *$G - L$ has a Λ-factor for every 3-vertex path L in G,*

(**z9**) $v(G) = 0 \bmod 6 \Rightarrow$ *for every 3-edge cut K of G and $S \subset K$, $|S| = 2$, there exists a Λ-factor P of G such that $E(P) \cap K = S$,*

(**t1**) $v(G) = 2 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exists $xy \in E(G)$ such that $G - \{x, y\}$ has a Λ-factor,*

(**t2**) $v(G) = 2 \bmod 6 \Rightarrow$ *$G - \{x, y\}$ has a Λ-factor for every $xy \in E(G)$,*

(**t3**) $v(G) = 2 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exists a 5-vertex path W such that x is the center vertex of W and $G - W$ has a Λ-factor* (see also **2.4** (a1.2) and Fig. 3),

(**t4**) $v(G) = 2 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ and $xy \in E(G)$ there exists a 5-vertex path W such that x is the center vertex of W, $xy \notin E(W)$, and $G - W$ has a Λ-factor* (see also **2.4** (a1.2) and Fig. 3),

(**f1**) $v(G) = 4 \bmod 6 \Rightarrow$ *$G - x$ has a Λ-factor for every $x \in V(G)$,*

(**f2**) $v(G) = 4 \bmod 6 \Rightarrow$ *$G - \{x, e\}$ has a Λ-factor for every $x \in V(G)$ and $e \in E(G)$,*

(**f3**) $v(G) = 4 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exists a 4-vertex path Z such that x is an inner vertex of Z and $G - Z$ has a Λ-factor* (see also **2.4** (a2.2) and Fig. 4),

(**f4**) $v(G) = 4 \bmod 6 \Rightarrow$ *for every $x \in V(G)$ there exists $xy \in E(G)$ and a 4-vertex path Z such that x is an inner vertex of Z, $xy \notin E(Z)$, and $G - Z$ has a Λ-factor* (see also **2.4** (a2.2) and Fig. 4),

(**f5**) $v(G) = 4 \bmod 6 \Rightarrow$ *for every $xy \in E(G)$ there exists a 4-vertex path Z such that xy is the middle edge of Z and $G - Z$ has a Λ-factor* (see also **2.4** (a2.2) and Fig. 4),

(**f6**) $v(G) = 4 \bmod 6 \Rightarrow$ *for every $z \in V(G)$ and every 3-vertex path xyz there exists a 4-vertex path Z such that $xyz \subset Z$, z is an end-vertex of Z, and $G - Z$ has a Λ-factor* (see also **2.4** (a2.2) and Fig. 4).

Theorem **3.1** follows from **3.6** – **3.28** below. In [11] we have shown that claims (z1) - (z5) are true for cubic, 3-connected, and claw-free graphs. In [9] we have proved the following results related to (t3), (t4), and (f3) - (f6).

3.2 *Let G be a cubic 2-connected graph. Then*

(a1) *if $G - x$ has a Λ-factor for some vertex x in G* (and so $v(G) = 4 \bmod 6$)*, then there exists a 4-vertex path Π in G such that $G - \Pi$ has a Λ-factor,*

(a2) *if $G - x, y$ has a Λ-factor for some edge xy in G* (and so $v(G) = 2 \bmod 6$)*, then there exists a 5-vertex path W and two disjoint 4-vertex paths Π_1 and Π_2 in G such that $G - W$ and $G - (\Pi_1 \cup \Pi_2)$ have Λ-factors, and*

(a3) *if $v(G) = 4 \bmod 6$ and G has a Hamiltonian cycle, then there exists a claw $Y = K_{1,3}$*

in G such that $G - Y$ has a Λ-factor.

From claim $(a3)$ in **3.2** it follows that if G is a cubic Hamiltonian graph with $v(G) = 4 \bmod 6$, then its domination number is at most $\lfloor v(G)/3 \rfloor$.

The remarks below show that if claims $(z7)$, $(z8)$, $(t2)$ - $(t4)$, and $(f1)$ - $(f6)$ in **3.1** are true, then they are best possible in some sense.

$(r1)$ Obviously claim $(z7)$ is not true if condition "$|X| = 2$" is replaced by condition "$|X| = 3$". Namely, if G is a cubic 3-connected graph, $v(G) = 0 \bmod 6$, X is a 3-edge cut in G, and the two components of $G - X$ have the number of vertices not divisible by 3, then clearly $G - X$ has no Λ-factor. Moreover, for every 3-edge graph Z with no isolated vertices and distinct from $K_{1,3}$, there exist infinitely many pairs (G, X) such that G is a cubic 3-connected graph, $X \subset E(G)$, $|X| = 3$, $G[X]$ (the subgraph of G induced by X is isomorphic to Z, and $G - X$ is connected and has no Λ-factor. Also in Section 4 we describe an infinite set of cubic 3-connected graphs G such that such that $G - E(T)$ has no Λ-factor for every triangle T in G.

$(r2)$ There exist infinitely many triples (G, L, e) such that G is a cubic, 3-connected, bipartite, and planar graph, $v(G) = 0 \bmod 6$, L is a 3-vertex path in G, $e \in E(G - L)$, and $(G - e) - L$ has no Λ-factor, and so claim $(z8)$ is tight. Indeed, let $v \in V(G)$, $N(v, G) = \{x, y, z\}$, and $yz \in E(G)$, and so vyz is a triangle. Since G is 3-connected, x is not adjacent to $\{y, z\}$. Let L be a 3-vertex path in $G - v$ containing yz. Then x is an isolated vertex in $(G - e) - L$, and therefore $(G - e) - L$ has no Λ-factor. Similarly, let $S = abcda$ be a 4-cycle in G, $e = xd$ be an edge in a vertex in $G - E(S)$, and $L = abc$ be a 3-vertex path in G. Then again x is an isolated vertex in $(G - e) - L$, and therefore $(G - e) - L$ has no Λ-factor.

$(r2')$ The arguments in $(r2)$ also show that there exist infinitely many triples (G, L, L') such that $G - (L \cup L')$ has no Λ-factor, where L and L' are disjoint 3-vertex paths in G, and G is a cubic, 3-connected, bipartite, and planar graph, and so claim $(z8)$ is tight in this sense as well. Indeed, they can be obtained from the constructions in $(r2)$ by replacing vertex x by a 3-vertex path in $G - L$ containing x.

In Section 6 (see **6.1**) we give a construction providing infinitely many cubic cyclically 6-connected graphs G such that on the one hand, $(G - e) - L$ has a Λ-factor for every 3-vertex path L in G and every edge in $G - L$, and on the other hand, G has pairs of disjoint 3-vertex paths L and L' such that $G - (L \cup L')$ has no Λ-factor.

$(r3)$ There exist infinitely many triples (G, xy, e) such that G is a cubic, 3-connected, bipartite, and planar graph, $v(G) = 2 \bmod 6$, $xy \in E(G)$, $e \in E(G - \{x, y\})$, and $G - \{x, y, e\}$ has no Λ-factor, and so claim $(t2)$ is tight.

$(r4)$ There exist infinitely many triples (G, x, y) such that G is a cubic 3-connected graph, $v(G) = 2 \bmod 6$, $\{x, y\} \subset V(G)$, $x \ne y$, $xy \notin E(G)$, and $G - \{x, y\}$ has no Λ-factor, and so claim $(t2)$ is not true if vertices x and y are not adjacent.

$(r5)$ There exist infinitely many quadruples (G, a, b, x) such that G is a cubic 3-

connected graph with no 3-cycles and no 4-cycles, $v(G) = 4 \bmod 6$, $x \in V(G)$, a and b are non-adjacent edges in $G - x$, and $G - \{x, a, b\}$ has no Λ-factor, and so claim $(f2)$ is tight.

$(r6)$ There exist infinitely many triples (G, L, x) such that G is a cubic 3-connected graph with no 3-cycles and no 4-cycles, $v(G) = 4 \bmod 6$, $x \in V(G)$, L is a 3-vertex path in $G - x$, and $G - \{x, L\}$ has no Λ-factor (see claim $(f1)$).

$(r7)$ There exist infinitely many pairs (G, Π) such that G is a cubic 3-connected graph, $v(G) = 4 \bmod 6$, Π is a 4-vertex path in G, and $G - \Pi$ has no Λ-factor (see claims $(f3)$ - $(f6)$).

$(r8)$ There exist infinitely many triples (G, W) such that G is a cubic 3-connected graph, $v(G) = 2 \bmod 6$, W is a 5-vertex path in G, and $G - W$ has no Λ-factor (see claims $(t3)$ and $(t4)$).

The above discussion in $(r2)$ leads to the following question.

3.3 Problem. *Is the following claim true ? Suppose that G is a cubic 3-connected graph with no triangles and no squares (or moreover, cyclically 5-connected), e is an edge in G, and L is a 3-vertex path in $G - e$. Then $(G - e) - L$ has a Λ-factor.*

We need the following two results obtained before.

3.4 [7] *Let* $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2) *and P be a Λ-factor of G. Suppose that each A^i is a cubic graph and $v(A^i) = 0 \bmod 6$. Then $cmp(P^i) \in \{1, 2\}$ for every $i \in \{1, 2, 3\}$.*

Proof. Let $i \in \{1, 2, 3\}$. Since D^i is a matching and P^i consists of the components of P meeting D^i, clearly $cmp(P^i) \le 3$. Since $v(A^i) = -1 \bmod 6$, we have $cmp(P^i) \ge 1$. It remains to show that $cmp(P^i) \le 2$. Suppose, on the contrary, that $cmp(P^1) = 3$. Since P is a Λ-factor of G and $v(A^1 - a^1) = -1 \bmod 6$, clearly $v(P^1) \cap V(A^1 - a^1) = 5$ and we can assume (because of symmetry) that P_1 consists of three components $a_3^1 z_3 a_3^2$, $z_1 a_1^1 y^1$, and $z_2 a_2^1 u^1$ for some $y^1, u^1 \in V(A^1)$ Then $cmp(P^3) = 0$, a contradiction. □

3.5 [7] *Let A be a graph, $e = aa_1 \in E(A)$, and $G = Y(A, a)$* (see Fig. 2)*. Suppose that*

$(h1)$ A is cubic,

$(h2)$ $v(A) = 0 \bmod 6$, and

$(h3)$ a has no Λ-factor containing $e = aa_1$.

Then $v(G) = 0 \bmod 6$ and G has no Λ-factor.

Proof (uses **3.4**). Suppose, on the contrary, that G has a Λ-factor P. By definition of $G = Y(A, a)$, each A^i is a copy of A and edge $e^i = a^i a^i_1$ in A^i is a copy of edge $e = aa_1$ in A By **3.4**, $cmp(P^i) \in \{1, 2\}$ for every $i \in \{1, 2, 3\}$. Since P is a Λ-factor of G and $v(A^i - a^i) = -1 \bmod 6$, clearly $E(P) \cap D^i$ is an edge subset of a Λ–factor of A^i for every

$i \in \{1,2,3\}$ (we assume that edge $z_j a^i_j$ in G is edge $a^i a^i_j$ in A^i). Since $a^1 a^i_1$ does not belong to any Λ-factor of A^i for every $i \in \{1,2,3\}$, clearly $E(P) \cap \{z_1 a^1_1, z_1 a^2_1, z_1 a^3_1\} = \emptyset$. Therefore $z_1 \not\in V(P)$, and so P is not a Λ-factor of G, a contradiction. □

3.6 $(z1) \Leftrightarrow (z2)$.

Proof (uses **2.3** and **3.4**). Obviously, $(z1) \Leftarrow (z2)$. We prove $(z1) \Rightarrow (z2)$.

Suppose, on the contrary, that $(z1)$ is true but $(z2)$ is not true, i.e., there is a cubic 3-connected graph A and $aa_1 \in E(G)$ such that $v(A) = 0 \bmod 6$ and every Λ-factor of G contains aa_1. Let $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$, where each (A^i, a^i) is a copy of (A, a) and edge $a^i a^i_1$ in A^i is a copy of edge aa_1 in A (see Fig. 2). Since A is cubic and 3-connected, by **2.3**, G is also cubic and 3-connected. Obviously, $v(G) = 0 \bmod 6$. By $(z1)$, G has a Λ-factor P. Since each $v(A^i) = 0 \bmod 6$, by **3.4**, $cmp(P^i) \in \{1,2\}$ for every $i \in \{1,2,3\}$. Since P is a Λ-factor of G and $v(A^i - a^i) = -1 \bmod 6$, clearly $E(P) \cap D^i$ is an edge subset of a Λ-factor of A^i for every $i \in \{1,2,3\}$ (we assume that edge $z_j a^i_j$ in G is edge $a^i a^i_j$ in A^i). Since $a^1 a^i_1$ belongs to every Λ-factor of A^i for every $i \in \{1,2,3\}$, clearly $\{z_1 a^1_1, z_1 a^2_1, z_1 a^3_1\} \subseteq E(P)$. Therefore vertex z_1 has degree three in P, and so P is not a Λ-factor of G, a contradiction. □

3.7 $(z1) \Leftrightarrow (z3)$.

Proof. Claim $(z1) \Leftarrow (z3)$ is obvious. Claim $(z1) \Rightarrow (z3)$ follows from **3.5**. □

3.8 $(z1) \Leftrightarrow (z4)$.

Proof (uses **2.2** and **2.4** $(a1)$). Obviously, $(z1) \Leftarrow (z4)$. We prove $(z1) \Rightarrow (z4)$.

Suppose that $(z1)$ is true but $(z4)$ is not true, i.e., there is a cubic 3-connected graph A and $a \in V(A)$ such that $v(A) = 0 \bmod 6$ and a has degree one in every Λ-factor of A. Let G be the graph obtained from $B = K_{3,3}$ by replacing each vertex v of B by a copy A_v of $A - a$ (see Fig. 5). Obviously, $v(G) = 0 \bmod 6$ and by **2.2**, G is a cubic 3-connected graph. By $(z1)$, G has a Λ-factor P. If $uv \in E(P)$, then let $L(uv)$ denote the component of P containing uv. Let $V(B) = \{1, \ldots, 6\}$.

Since vertex a has degree one in every Λ-factor of A and each (A^v, a^v) is a copy of (A, a), by **2.4** $(a1)$, we have $cmp(P^v) \in \{1,3\}$.

(**p1**) Suppose that there is $v \in V(B)$ such that $cmp(P^v) = 3$. By symmetry of B, we can assume that $v = 1$ and, by **2.4** $(a1.3)$, $1 \neg 4$, $6 \neg 1$, and $2 \neg 1$ (see Fig. 5). Let $x \in \{2,6\}$. Since $cmp(P^x) \in \{1,3\}$ and $|V(L(1x) \cap V(A_x)| = 1$, clearly $cmp(P^x) = 3$ and, by **2.4** $(a1.3)$, $5 \neg x$, $3 \neg x$. Then by **2.4** $(a1.2)$, $cmp(P^s) = 2$ for $s \in \{3,5\}$, a contradiction.

(**p2**) Now suppose that $cmp(P^v) = 1$ for every $v \in V(B)$. By symmetry, we can assume $1 \neg 2$. Then P^2 contradicts **2.4** $(a1.1)$. □

3.9 $(z1) \Leftrightarrow (t1)$.



Figure 5: $(z1) \Rightarrow (z4)$

Proof (uses **2.3**). If $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2) and each A^i is cubic and 3-connected, then by **2.3**, G is also cubic and 3-connected.

(**p1**) We first prove $(z1) \Rightarrow (t1)$. Suppose, on the contrary, that $(z1)$ is true but $(t1)$ is not true, i.e., there is a cubic 3-connected graph A and $a \in V(A)$ such that $v(A) = 2 \bmod 6$ and $A - \{a, y\}$ has no Λ-factor for every vertex y in A adjacent to a. Let each (A^i, a^i) above be a copy of (A, a), and so $A^i - \{a^i, a^i_j\}$ has no Λ-factor for every $i, j \in \{1, 2, 3\}$. Obviously $v(G) = 0 \bmod 6$. By $(z1)$, G has a Λ-factor P. Then it is easy to see that since each $v(A^i - a^i) = 1 \bmod 6$, there are $r, s, j \in \{1, 2, 3\}$ such that $r \neq s$ and $P^s = a^s_j z_j a^r_j$. Since P is a Λ-factor of G, clearly $P \cap (A^s - \{z_j, a^s_j\})$ is a Λ-factor of $A^s - \{z_j, a^s\} = A^s - \{a^s, a^s_j\}$, a contradiction.

(**p2**) Now we prove $(z1) \Leftarrow (t1)$. Suppose, on the contrary, that $(t1)$ is true but $(z1)$ is not true, i.e., there is a cubic 3-connected graph A such that $v(A) = 0 \bmod 6$ and A has no Λ-factor.

Let (A^i, a^i) be a copy of (A, a) for $i \in \{1, 2\}$, where $a \in V(A)$, and (A^3, a^3) be a copy of (H, h) for some cubic 3-connected graph H and $h \in V(H)$, where $v(H) = 2 \bmod 6$. Obviously $v(G) = 2 \bmod 6$. Suppose that P is a Λ-factor of $G - \{z_1 a^3_1\}$. Then $cmp(P^1) \leq 2$. Since $v(A^1 - a^1) = -1 \bmod 6$, we have $cmp(P^1) \geq 1$. Now since P is a Λ–factor of $G - \{z_1 a^3_1\}$ and $v(A^1 - a^1) = -1 \bmod 6$, clearly $E(P) \cap D^1$ is an edge subset of a component of a Λ-factor of A^1 (we assume that edge $z_j a^1_j$ in G is edge $a^1 a^1_j$ in A^1). Therefore A has a Λ-factor, a contradiction. □

3.10 $(z1) \Leftrightarrow (f1)$.

Proof (uses **2.3**). By **2.3**, if $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2) and each A^i is cubic and 3-connected, then G is also cubic and 3-connected.

(**p1**) We first prove $(z1) \Rightarrow (f1)$. Suppose, on the contrary, that $(z1)$ is true but $(f1)$ is not true, i.e., there is a cubic 3-connected graph A such that $v(A) = 4 \bmod 6$ and $A - a$ has no Λ-factor for some $a \in V(A)$. Let each (A^i, a^i) above be a copy of (A, a). Obviously, $v(G) = 0 \bmod 6$. By $(z1)$, G has a Λ-factor P. Let P^i be the union of components in P meeting D^i and $E^i = E^i(P)$.

Since P is a Λ-factor of G and $v(A^i - a^i) = 3 \bmod 6$, clearly each $d(z_j, P) \le 2$ and each $|E^i| \in \{0, 2, 3\}$. Since $A^i - a^i$ has no Λ-factor, each $|E^i| \in \{2, 3\}$, and so each $d(z_j, P) \ge 2$. Thus, each $d(z_j, P) = 2$, and so each z_j is the center of a 3-vertex path in P and each $|E^i| = 2$. Now, since $v(A^i - a^i) = 3 \bmod 6$, $P \cap (A^i - a^i - V(P^i))$ is not a Λ-factor of $A^i - a^i - V(P^i))$. Therefore P is not a Λ-factor of G, a contradiction.

(**p2**) Now we prove $(z1) \Leftarrow (f1)$. Suppose, on the contrary, that $(f1)$ is true but $(z1)$ is not true, i.e., there is a cubic 3-connected graph A such that $v(A) = 0 \bmod 6$ and A has no Λ-factor. Let (A^i, a^i) above be a copy of (A, a) for $i \in \{1, 2\}$ and some $a \in V(A)$ and (A^3, a^3) is a copy of (H, h) for some cubic 3-connected graph H and $h \in V(H)$, where $v(H) = 4 \bmod 6$ (see Fig. 2). Then $v(G) = 4 \bmod 6$. Let $x \in V(H - h)$. Suppose that $G - x$ has a Λ-factor P. Since A has no Λ-factor, clearly $|E^1(P)| = |E^2(P)| = 3$. Then P is not a Λ-factor of $G - x$, and so $(f1)$ is not true, a contradiction. □

3.11 $(z4) \Leftrightarrow (t2)$.

Proof (uses **2.3**, **3.8**, and **3.9**). Let $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2). By **2.3**, if each A^i is cubic and 3-connected, then G is also cubic and 3-connected.

(**p1**) We first prove $(z4) \Leftarrow (t2)$. Obviously $(t2) \Rightarrow (t1)$. By **3.8**, $(z1) \Leftrightarrow (z4)$. By **3.9**, $(z1) \Leftrightarrow (t1)$. The result follows.

(**p2**) Now we prove $(z4) \Rightarrow (t2)$. Suppose, on the contrary, that $(z4)$ is true but $(t2)$ is not true. Then there exists a cubic 3-connected graph A and $aa_1 \in E(G)$ such that $v(A) = 2 \bmod 6$ and $A - \{a, a_1\}$ has no Λ-factor. Let each (A^i, a^i) above be a copy of (A, a) and edge $a^i a^i_1$ in A^i be a copy of edge aa_1 in G. Obviously $v(G) = 0 \bmod 6$. Let $L^i = a^j_1 z_1 a^k_1$, where $\{i, j, k\} = \{1, 2, 3\}$. By $(z4)$, G has a Λ-factor P containing L^i for some $i \in \{1, 2, 3\}$, say for $i = 3$. If $s \in \{1, 2\}$, then $cmp(P^s) = 3$, because $A^s - \{a^s, a^s_1\}$ has no Λ-factor and $v(A^s - a^s) = 1 \bmod 6$. Also $cmp(P^3) \ge 1$, because $v(A^3 - a^3) = 1 \bmod 6$. Then $P^1 \cup P^2 \cup P^3$ has at least four components each meeting $\{z_1, z_2, z_3\}$, a contradiction. □

3.12 $(z2) \Rightarrow (z5)$.

Proof (uses **2.1** and **2.4** $(a1)$). Suppose, on the contrary, that $(z2)$ is true but $(z5)$ is not true. Then there exists a cubic 3-connected graph A and $a \in V(A)$ such that at most one 3-vertex path, centered at a belongs to a Λ-factor of A. It is sufficient to prove



Figure 6: $(z2) \Rightarrow (z5)$

our claim in case when A has exactly one 3-vertex path, say $L = a_1 a a_2$, centered at a and belonging to a Λ-factor of A. Let $e_i = aa_i$, and so $E(L) = \{e_1, e_2\}$.

Let B be the graph-skeleton of the three-prism: $V(B) = \{1, \ldots, 6\}$ and B is obtained from two disjoint triangles 123 and 456 by adding three new edges 14, 25, and 36.

Let each (A^v, a^v, a_1^v, a_2^v), $v \in V(B)$ be a copy of (A, a, a_1, a_2), and so edge $e_i^v = a^v a_i^v$ in A^v is a copy of edge $e_i = aa_i$ in A, $i \in \{1, 2\}$. We also assume that all A^v's are disjoint. Let G be a graph obtained from B by replacing each $v \in V(B)$ by $A_v = A^v - a^v$ (see Fig. 6). We recall the following notation from Section 2:

(1) $E'(G) = E(G) \setminus \cup\{E(A_v) : v \in V(B)\}$,

(2) $\alpha : E(B) \to E'(G)$ is the bijection such that if $uv \in E(B)$, then $\alpha(uv)$ is an edge in G having one end-vertex in A_u and the other in A_v, and

(3) if P is a Λ-packing in G, then for $uv \in E(B)$, $u \neq v$, we write $u \neg^P v$ (or simply, $u \neg v$), if P has a 3-vertex path L such that $\alpha(uv) \in E(L)$ and $|V(A_u) \cap V(L)| = 1$.

Given $v \in V(B)$, let $S(v)$ be the set of two edges e_i' in $E'(G)$ such that edge e_i' is incident to vertex a_i^v in G, $i \in \{1, 2\}$. We assume that each vertex v in B is replaced by A_v (to obtain G) in such a way that
$S(1) = \{\alpha(13), \alpha(14\}$, $S(2) = \{\alpha(21), \alpha(25)\}$, $S(3) = \{\alpha(32), \alpha(36)\}$,
$S(4) = \{\alpha(45), \alpha(46)\}$, $S(5) = \{\alpha(54), \alpha(56)\}$, $S(6) = \{\alpha(63), \alpha(64)\}$.
In Figure 6 the edges in $S(v)$ are marked for every $v \in V(B)$.

By **2.1**, G is a cubic 3-connected graph. Since $v(B) = 0 \bmod 6$, clearly also $v(G) = 0 \bmod 6$. By $(z2)$, $G' = G - \alpha(36)$ has a Λ-factor P.

We know that A has exactly one 3-vertex path $L = a_1 a a_2$ centered at a and belonging to a Λ-factor of A and that each (A^v, a^v, a_1^v, a_2^v) is a copy of (A, a, a_1, a_2), and so $v(A^v -$

$a^v) = -1 \bmod 6$. Therefore, by **2.4** ($a1$) and the assumption on A, the Λ-factor P satisfies the following condition for every $v \in V(B)$:

c(**v**) if $cmp(P^v) = 2$, then $v\neg a$ and $v\neg b$, where $\{\alpha(va), \alpha(vb)\} = S(v)$.

Obviously, $|D^3| = |D^6| = 2$ in $G - \alpha(36)$. Therefore $cmp(P^3) \le 2$ and $cmp(P^6) \le 2$. Since $\alpha(36) \in S(3) \cap S(6)$, by conditions **c**(**3**) and **c**(**6**), $cmp(P^3) = cmp(P^6) = 1$. Now by **2.4** ($a1$), $x'\neg 3$ for some $x' \in \{1, 2\}$ and $y'\neg 6$ for some $y' \in \{4, 5\}$.

(**p1**) Suppose that $1\neg 3$. Assume first that $\alpha(14) \notin E(P)$. Then $cmp(P^1) \le 2$. By **2.4** ($a1$), $cmp(P^1) = 2$. This contradicts **c**(**1**). Thus we must have $\alpha(14) \in E(P)$.

(**p1**.**1**) Suppose that $4\neg 6$.
Suppose that $1\neg 4$. Then by **2.4** ($a1$), $5\neg 4$ and $5\neg 2$. This contradicts **c**(**5**).
Now suppose that $4\neg 1$. This contradicts **c**(**4**).

(**p1**.**2**) Suppose that $5\neg 6$. Then $cmp(P^4) \le 2$.
Suppose that $1\neg 4$. By **2.4** ($a1.1$), $cmp(P^4) = 1$. Then $5\neg 2$, which contradicts **c**(**5**).
Now suppose that $4\neg 1$. Then $cmp(P^4) = 2$, which contradicts **c**(**4**).

(**p2**) Now suppose that $2\neg 3$. Then $\alpha(13) \notin E(P)$, and therefore $cmp(P^1) \le 2$. By **c**(**2**) and **2.4** ($a1.3$), $1\neg 2$ (and $5\neg 2$). Then by **2.4** ($a1.2$), $cmp(P^1) = 2$, which contradicts **c**(**1**). □

Since ($z7$) ⇒ ($z2$), we obtain from **3.12**:

3.13 ($z7$) ⇒ ($z5$).

3.14 ($z1$) ⇔ ($z6$).

Proof. Obviously, ($z1$) ⇐ ($z6$) and ($z5$) ⇒ ($z6$). As to ($z1$) ⇒ ($z6$), it follows from **3.6** and **3.12**. □

3.15 ($z7$) ⇒ ($z8$).

Proof 1. Suppose, on the contrary, that ($z7$) is true but ($z8$) is not true. Then there exists a cubic 3-connected graph A and a 3-path $L = a_1 a a_2$ in A such that $v(A) = 0 \bmod 6$ and $A - L$ has no Λ-factor. Let $N(a, A) = \{a_1, a_2, a_3\}$. Let $(A^i; a^i, a^i_1, a^i_2, a^i_3)$, $i \in \{1, 2\}$, be two copies of $(A; a, a_1, a_2, a_3)$ and A^1, A^2 be disjoint graphs, and so $L^i = a^i_1 a^i a^i_2$ in A^i is a copy of $L = a_1 a a_2$ in A. Let $H = A^1 a^1 \sigma a^2 A^2$, where $\sigma : N(a^1, A^1) \to N(a^2, A^2)$ is a bijection such that $\sigma(a^1_i) = a^2_i$ for $i \in \{1, 2, 3\}$. Let G be the graph obtained from H by subdividing edge $a^1_j a^2_j$ by a new vertex v_j for $j \in \{1, 3\}$ and by adding a new edge $v_1 v_3$. Obviously, G is a cubic 3-connected graph and $v(G) = 0 \bmod 6$. By ($z7$), $G' = G - \{a^1_3 v_3, a^2_3 v_3\}$ has a Λ-factor P. Since $d(v_3, G') = 1$, P has a 3-vertex path $L_1 = v v_1 v_3$. By symmetry of G, we can assume that $v = a^1_1$. Since $v(A^1 - a^1) = -1 \bmod 6$, P has a 3-vertex path $L_2 = a^1_2 a^2_2 b$ for some $b \in V(A^2 - a^2)$. Then $P \cap (A^1 - L)$ is a Λ-factor of $A^1 - L$, a contradiction. □



Figure 7: $(z7) \Rightarrow (z8)$

Proof 2 (uses **2.4** $(a1)$). Suppose, on the contrary, that $(z7)$ is true but $(z8)$ is not true. Then there is a cubic 3-connected graph A and a 3-path $L = a_1aa_2$ in A such that $v(A) = 0 \bmod 6$ and $A - L$ has no Λ-factor. Let $N(a, A) = \{a_1, a_2, a_3\}$.

Let B, $\{(A^v, a^v, a_1^v, a_2^v, a_3^v) : v \in V(B)\}$, and G be as in **3.12** (see Fig. 7). Given $v \in V(B)$, let $S(v)$ be the set of two edges e_i' in $E'(G)$ such that edge e_i' is incident to vertex a_i^v in G, $i \in \{1, 2\}$. We assume that each vertex v in B is replaced by A^v (to obtain G) in such a way that

$S(1) = \{\alpha(12), \alpha(13)\}$, $S(2) = \{\alpha(21), \alpha(23)\}$, $S(3) = \{\alpha(32), \alpha(31)\}$,
$S(4) = \{\alpha(45), \alpha(46)\}$, $S(5) = \{\alpha(54), \alpha(56)\}$, $S(6) = \{\alpha(64), \alpha(65)\}$.

In Figure 7 the edges in $S(v)$ are marked for every $v \in V(B)$.

Since $v(G) = 0 \bmod 6$ and G is cubic and 3-connected, by $(z7)$, $G - \{\alpha(14), \alpha(36)\}$ has a Λ-factor P. By **2.4** $(a1)$ and the lack of Λ-factor in each of $A^v - a_1^v a^v a_2^v$, the Λ-factor P satisfies the following condition for every $v \in V(B)$:

c(v) if $cmp(P^v) = 2$, then $v \neg a$ and $v \neg b$, where $\{\alpha(va), \alpha(vb)\} \neq S(v)$.

Obviously, $|D^i| = 2$ in $G - \{\alpha(14), \alpha(36)\}$, and so $cmp(P^i) \le 2$ for $i \in \{1, 3\}$. Since $S(1) = \{12, 13\}$ and $S(3) = \{31, 32\}$, by conditions **c(1)** and **c(4)** we have $cmp(P^1) = cmp(P^3) = 1$. Now by **2.4** $(a1.1)$, $2 \neg 1$ and $2 \neg 3$. This contradicts **c(2)**. □

3.16 $(t2) \Leftrightarrow (z7)$.

Proof (uses **3.8** and **3.11**). We first prove $(t2) \Rightarrow (z7)$. Let G be a cubic, 3-connected graph with $v(G) = 0 \bmod 6$ and $a = a_1a_2$, $b = b_1b_2$ be two distinct edges of G. Let G' be the graph obtained from G as follows: subdivide edge a_1a_2 by a new vertex a' and edge b_1b_2 by a new vertex b' and add a new edge $e = a'b'$. Then G' is a cubic and 3-connected graph, $v(G') = 2 \bmod 6$, and $G - \{a, b\} = G' - \{a', b'\}$. By $(t2)$, $G' - \{a', b'\}$ has a Λ-factor.

Now we prove $(t2) \Leftarrow (z7)$. Obviously, $(z7) \Rightarrow (z1)$. By **3.8**, $(z1) \Rightarrow (z4)$ and by **3.11**, $(z4) \Rightarrow (t2)$. Implication $(t2) \Leftarrow (z7)$ also follows from obvious $(z8) \Rightarrow (z4)$, from $(z7) \Rightarrow (z8)$, (**3.15**), and from $(z4) \Rightarrow (t2)$ (**3.11**). $\square$

Here is a direct proof of $(z7) \Rightarrow (t2)$.

3.17 $(z7) \Rightarrow (t2)$.

Proof. Let G be a cubic, 3-connected graph, $v(G) = 2 \bmod 6$, $xy \in E(G)$, $N(x, G) = \{x_1, x_2, y\}$, and $N(y, G) = \{y_1, y_2, x\}$. Let $G_1 = G - \{x, y\} \cup E_1$, $G_2 = G - \{x, y\} \cup E_2$, and $G_3 = G - \{x, y\} \cup E_3$, where $E_1 = \{x_1y_1, x_2y_2\}$, $E_2 = \{x_1y_2, x_2y_1\}$, and $E_3 = \{x_1x_2, y_1y_2\}$. Obviously, each G_i is a cubic graph and $v(G_i) = 0 \bmod 6$. It is easy to see that since G is 3-connected, there is $s \in \{1, 2, 3\}$ such that G_s is 3–connected. Clearly $G_s - E_s = G - \{x, y\}$. By $(z7)$, $G_s - E_s$ has a Λ-factor. $\square$

3.18 $(z8) \Leftrightarrow (z7)$.

Proof. Obviously, $(z8) \Rightarrow (z4)$. By **3.11**, $(z4) \Rightarrow (t2)$ and by **3.16**, $(t2) \Rightarrow (z7)$. Therefore $(z8) \Rightarrow (z7)$. By **3.15**, $(z7) \Rightarrow (z8)$. $\square$

3.19 $(z1) \Leftrightarrow (z8)$.

Proof. Obviously, $(z8) \Rightarrow (z1)$. By **3.8**, $(z1) \Rightarrow (z4)$. By **3.11**, $(z4) \Rightarrow (t2)$. By **3.16**, $(t2) \Rightarrow (z7)$. By **3.15**, $(z7) \Rightarrow (z8)$. Therefore $(z1) \Rightarrow (z8)$. $\square$

3.20 $(z8) \Rightarrow (f1)$.

Proof. Let G be a cubic 3-connected graph, $v(G) = 4 \bmod 6$, $x \in V(G)$, and $N(x, G) = \{x_1, x_2, x_3\}$. Let G' be the graph obtained from G by replacing x by a triangle T with $V(T) = \{x'_1, x'_2, x'_3\}$ so that $x_ix'_i \in E(G')$, $i \in \{1, 2, 3\}$. Since $v(G) = 4 \bmod 6$, clearly $v(G') = 0 \bmod 6$. Consider the 3-vertex path $L' = x'_1x'_2x'_3$ in G'. By $(z8)$, $G' - L'$ has a Λ-factor P'. Obviously, $P' - L'$ is a Λ-factor of $G' - L'$ and $G - x = G' - L'$. $\square$

3.21 $(z8) \Rightarrow (f2)$.

Proof (uses **3.20**). Let G be a cubic 3-connected graph, $v(G) = 4 \bmod 6$, $x \in V(G)$, and $e = y_1y_2 \in E(G)$. We want to prove that if $(z8)$ is true, then $G - \{x, e\}$ has a Λ-factor. If $x \in \{y_1, y_2\}$, then $G - \{x, e\} = G - x$, and therefore, by **3.20**, our claim is true. So we assume that $x \notin \{y_1, y_2\}$. Let $N(x, G) = \{x_1, x_2, x_3\}$. Let G' be the graph obtained from G by subdividing edge y_1y_2 by a vertex y and edge xx_3 by a vertex z and by adding a new edge yz (see Fig. 8). Since $v(G) = 4 \bmod 6$, clearly $v(G') = 0 \bmod 6$. Since $x \notin \{y_1, y_2\}$ and G is cubic and 3-connected, G' is also cubic and 3-connected. Obviously, $L = xzy$ is a 3-vertex path in G' and $G - \{x, e\} = G - \{x, y_1y_2\} = G' - L$. By $(z8)$, $G' - L$ has a Λ-factor. $\square$



Figure 8: $(z8) \Rightarrow (f2)$

3.22 $(f2) \Rightarrow (t4)$.

Proof. Let G be a cubic, 3-connected graph, $v(G) = 2 \bmod 6$, $x \in V(G)$, and $N(x, G) = \{x_1, x_2, x_3\}$. Let G' be the graph obtained from G by replacing x by a triangle T with $V(T) = \{x'_1, x'_2, x'_3\}$ so that $x_i x'_i \in E(G')$, $i \in \{1, 2, 3\}$. Since $v(G) = 2 \bmod 6$, clearly $v(G') = 4 \bmod 6$. By $(f2)$, $G' - \{x'_i, x'_j x'_k\}$ has a Λ-factor P_i, where $\{i, j, k\} = \{1, 2, 3\}$. Since $x_j x'_j$ and $x_k x'_k$ are dangling edges in $G' - \{x'_i, x'_j x'_k\}$, clearly $x_j x'_j, x_k x'_k \in E(P_i)$ and $d(x'_j, P_i) = d(x'_k, P_i) = 1$. Let L_j and L_k be the components of P_i containing $x_j x'_j$ and $x_k x'_k$, respectively. Then $E(L_j) \cup E(L_k)$ induces in G a 5-vertex path W_i such that x is the center vertex of W_i and $x_i x'_i \notin E(W_i)$. $\square$

3.23 $(t3) \Rightarrow (z1)$.

Proof (uses **2.4** $(a1.3)$). Suppose, on the contrary, that $(t3)$ is true but $(z1)$ is not true, i.e., there is a cubic 3-connected graph A such that $v(A) = 0 \bmod 6$ and A has no Λ-factor. Let $a \in V(A)$. Let (A^i, a^i) above be a copy of (A, a) for $i \in \{1, 2\}$ and let (A^3, a^3) be such that $v(A^3) = 2 \bmod 6$. Let $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2). Obviously, $v(G) = 2 \bmod 6$. By $(t3)$, G has a 5-vertex path W such that z_3 is the center vertex of W and $G - W$ has a Λ-factor P. Obviously, $(A^i - a^i) \cap W = \emptyset$ for some $i \in \{1, 2, 3\}$.

Suppose that $(A^3 - a^3) \cap W = \emptyset$. Then W has an end-edge in $A^1 - a^1$ and in $A^2 - a^2$. Since A has no Λ-factor, by **2.4** $(a1.3)$, $D^i - e(W) \subseteq E(P)$ for $i \in \{1, 2\}$. Then P is not a Λ-factor of $G - W$, a contradiction.

Now suppose that $(A^3 - a^3) \cap W \neq \emptyset$. By symmetry, we can assume that $(A^2 - a^2) \cap W = \emptyset$. Then W has an end-edge in $A^1 - a^1$ and in $A^3 - a^3$. Then by **2.4** $(a1.3)$, $Cmp(P^1) = \{L_1, L_2\}$, where L_1 has an end-vertex in $A^1 - a^1$ and L_2 has an end-edge in $A^1 - a^1$. By symmetry, we can assume that $a^1_i z_i \in E(L_i)$ for $i \in \{1, 2\}$. Then $L_1 = a^1_1 z_1 y$, where $y \in \{a^2_1, a^3_1\}$ and z_2 is of degree one in P. Then, taking into account the values of $v((A^i - a^i)$ $i \in \{1, 2, 3\}$, P is not a Λ-factor of $G - W$, a contradiction. $\square$

3.24 $(z8) \Rightarrow (f4)$.

Proof. Let G be a cubic 3-connected graph, $v(G) = 4 \bmod 6$, $x \in V(G)$, and $N(x, G) = \{x_1, x_2, x_3\}$. Let G' be the graph obtained from G by replacing x by a triangle Δ with $V(\Delta) = \{x'_1, x'_2, x'_3\}$, so that $x_i x'_i \in E(G')$, $i \in \{1, 2, 3\}$. Since $v(G) = 4 \bmod 6$, clearly $v(G') = 0 \bmod 6$. Since G is cubic and 3-connected, G' is also cubic and 3-connected. Consider 3-vertex path $L'_i = x_j x'_j x'_k$ in G', where $\{i, j, k\} = \{1, 2, 3\}$. By $(z8)$, $G' - L'_i$ has a Λ-factor P'_i. Since $x_i x'_i$ is a dangling edge in $G' - L'_i$, clearly $x_i x'_i \in E(P'_i)$ and $d(x'_i, P_i) = 1$. Let Λ'_i be the 3-vertex path in P'_i containing $x'_i x_i$, say $\Lambda'_i = x'_i x_i z$. Consider in G the 4-vertex path $Z_k = x_k x x_i z$. Then x is an inner vertex of Z_k, $x x_k \notin E(Z_k)$, and $P'_i - \{L'_i \cup \Lambda'_i\}$ is a Λ-factor in $G - Z_k$. □

Let H' be a tree such that $V(H') = \{x, y\} \cup \{b^j : j \in \{1, 2, 3, 4\}$ and $E(H') = \{xy, b^1 x, b^2 x, b^3 y, b^4 y\}$. Let H_i, $i \in \{1, 2, 3\}$ be three disjoint copies of H' with $V(H_i) = \{x_i, y_i\} \cup \{b^j_i : j \in \{1, 2, 3, 4\}\}$. Let H be the graph obtained from these three copies by identifying for every j three vertices b^j_1, b^j_2, b^j_3 with a new vertex z^j. Let A^i, $i \in \{1, 2, 3, 4\}$, be a cubic graph, $a^i \in V(A^i)$, and let $G = H(A^1, a^1; A^2, a^2; A^3, a^3; A^4, a^4)$ be the graph obtained from H by replacing each z^j by $A^j - a^j$ assuming that all A^i's are disjoint.

3.25 $(f3) \Rightarrow (z1)$.

Proof (uses **2.2**). Suppose, on the contrary, that $(f3)$ is true but $(z1)$ is not true, i.e., there exists a cubic 3-connected graph A such that $v(A) = 0 \bmod 6$ and A has no Λ-factor. Let $a \in V(A)$. Let A^4 be any cubic 3-connected graph with $v(A^4) = 2 \bmod 6$ and each (A^i, a^i), $i \in \{1, 2, 3\}$, be a copy of (A, a).

Let the graph-composition $G = H(A^1, a^1; A^2, a^2; A^3, a^3; A^4, a^4)$ and its frame H be as defined above. Then H is a cubic 3-connected graph. Since each brick A^i is cubic and 3-connected, by **2.2**, G is also cubic and 3-connected. Obviously, $v(G) = 4 \bmod 6$. It is easy to see that $G - Z$ has no Λ-factor for every 4-vertex path Z in G such that y_1 is an inner vertex of Z. This contradicts $(f3)$. □

Obviously, $(f4) \Rightarrow (f3)$. Therefore from **3.25** we have $(f4) \Rightarrow (z1)$. Below we give a direct proof of this implication.

3.26 $(f4) \Rightarrow (z1)$.

Proof. Suppose, on the contrary, that $(f4)$ is true but $(z1)$ is not true, i.e., there exists a cubic 3-connected graph A such that $v(A) = 0 \bmod 6$ and A has no Λ-factor. Let $a \in V(A)$. Let (A^3, a^3) be a copy of (A, a) and let (A^i, a^i) for $i \in \{1, 2\}$ be copies of (B, b), where B is a cubic 3-connected graph, $v(B) = 2 \bmod 6$, and $b \in V(B)$. Let $G = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2). Obviously, $v(G) = 4 \bmod 6$. By $(f4)$, G has a 4-vertex path Z such that z_3 is an inner vertex of Z, $(A^3 - a^3) \cap Z = \emptyset$, and $G - Z$ has a Λ-factor, say P. Since A^3 has no Λ-factor, clearly P is not a Λ-factor of $G - Z$, a contradiction. □

Implication $(z8) \Rightarrow (f1)$ follows from obvious $(z8) \Rightarrow (z1)$ and from $(z1) \Rightarrow (f1)$, by

3.10. It also follows from obvious $(f2) \Rightarrow (f1)$ and from $(z8) \Rightarrow (f2)$, by **3.21**. Below we give a direct proof of this implication.

3.27 $(f6) \Rightarrow (f5) \Rightarrow (f4) \Rightarrow (z1) \Rightarrow (f6)$.

Proof (uses **3.26** and **3.19**). Obviously, $(f6) \Rightarrow (f5) \Rightarrow (f4)$. By **3.26**, $(f4) \Rightarrow (z1)$. Therefore $(f6) \Rightarrow (z1)$. It remains to prove $(z1) \Rightarrow (f6)$. By **3.19**, $(z1) \Rightarrow (z8)$. Thus it is sufficient to show that $(z8) \Rightarrow (f6)$. Let G be a cubic 3-connected graph, $v(G) = 4 \bmod 6$, and xyz is a 3-vertex path in G. Let $N(y, G) = \{x, z, s\}$ and G' be obtained from G by subdividing edges yz and ys by new vertices z' and s', respectively, and by adding a new edge $s'z'$. Then G' is a cubic 3-connected graph and $v(G') = 0 \bmod 6$. Consider the 3-vertex path $L' = zz's'$ in G'. By $(z8)$, G' has a Λ-factor P' containing L'. Since vertex y has degree one in $G' - L'$, clearly P' has a 3-vertex path $Q' = yxq$. Let Z be the 4-vertex path $qxyz$ in G. Then $P' - (L' \cup Q')$ is a Λ-factor in $G - Z$. □

3.28 $(z9) \Leftrightarrow (z1)$.

Proof. Obviously, $(z9) \Rightarrow (z8)$. By the above claims, $(z1)$, $(z8)$, $(t4)$, and $(f6)$ are equivalent. So we can use these claims to prove $(z9)$. Let G be a cubic 3-connected graph, K a 3-edge cut of G, $S \subset K$ and $|S| = 2$, and $v(G) = 0 \bmod 6$. If the edges of K are incident to the same vertex x in G (i.e., $K = D(x, G)$), then by $(z8)$, G has a Λ-factor P of G such that $E(P) \cap K = S$, and so our claim is true. So we assume that the edges in K are not incident to the same vertex in G. Then, since G is cubic and 3-connected, clearly 3-edge cut K forms a matching. Let A and B be the two component of $G - K$. By the above arguments, we have $v(A) \neq 1$ and $v(B) \neq 1$. Let A^b be the graph obtained from G by identifying the vertices of B with a new vertex b and similarly, let B^a be the graph obtained from G by identifying the vertices of A with a new vertex a, and so $A = A^b - b$ and $B = B^a - a$. Then $D(b, A^b) = D(a, B^a) = D$, and $S \subset D$. Let $S = \{a_1b_1, a_2b_2\}$, where $\{a_1, a_2\} \subset V(A)$ and $\{b_1, b_2\} \subset V(B)$. Obviously, S forms a 3-vertex path $S_A = a_1ba_2$ in A^b and a 3-vertex path $S_B = b_1ab_2$ in B^a. Since G is cubic and 3-connected, both A^b and B^a are also cubic and 3-connected. Since $v(G) = 0 \bmod 6$, there are two possibilities (up to symmetry):

$(c1)$ $v(A^b) = 0 \bmod 6$ and $v(B^a) = 2 \bmod 6$ and
$(c2)$ $v(A^b) = v(B^a) = 4 \bmod 6$.

Consider case $(c1)$. By $(z8)$, $A^b - S_A$ has a Λ-factor P_A. By $(t4)$, B^a has a 5-vertex path W such a is the center vertex of W, $S_B \subset W$, and $B^a - W$ has a Λ-factor P_B. Then $E(P_A) \cup E(P_B) \cup S$ induces a Λ-factor P in G such that $E(P) \cap K = S$.

Now consider case $(c2)$. By $(f6)$, we have:

(a) A^b has a 4-vertex path Z_A such that a_1 is an end-vertex of Z_A, $S_A \subset Z_A$, and $A^b - Z_A$ has a Λ-factor, P_A;

(b) likewise, B^a has a 4-vertex path Z_B such that b_2 is an end-vertex of Z_B, $S_B \subset Z_B$, and $B^a - Z_B$ has a Λ-factor P_B.

Then $E(P_A) \cup E(P_B) \cup S$ induces a Λ-factor P in G such that $E(P) \cap K = S$. $\square$

4 Properties of Λ-factors in graph-compositions

In this section we consider graph-compositions having Λ-factors and with all bricks having the same number of vertices modulo 6. It turns out that the Λ-factors of such graph-compositions intersect its 3-blockades in a special way and correspond to some special subgraphs in the frame of the composition.

Let $G = B\{(A^v, a^v) : v \in V(B)\}$ and P be a Λ-factor of G if any exists. Let $F(P,G)$ be the subgraph in B induced by the edge set $\alpha^{-1}(E'(G) \cap E(P))$. Let $\vec{F}(P,G)$ be the directed graph obtained from $F(P,G)$ by directing each edge uv in $F(P,G)$ from u to v, if $u \neg^p v$. For a Λ-factor P of B, let $\vec{P}$ denote the directed graph obtained from P by directing the two edges of each 3-vertex path L in P to the central vertex of L. We call $\vec{P}$ a Λ-difactor of B.

A subgraph C of a graph H is called a *cycle-packing* in H if each component of C is a cycle. For a cycle-packing Q in B, let $\vec{Q}$ denote the directed graph obtained from Q by directing its edges so that every cycle in $\vec{Q}$ becomes a directed cycle. We call $\vec{Q}$ a *cycle-dipacking* of B.

4.1 *Let B be a cubic 3-connected graph with no 3-blockades. Let G be the graph obtained from B by replacing each vertex by a triangle* (i.e., $G = B\{K_4, x\}$, where $x \in V(K_4)$, and so G is a claw-free graph).

$(a1)$ *Let P be a Λ-factor of G. Then*

- $(a1.1)$ *$\vec{F}(P,G)$ is a cycle-dipacking in B,*
- $(a1.2)$ *$|K \cap E(P)| \in \{0,2\}$ for every 3-blockade K of G, and*
- $(a1.3)$ *$E(T) \cap E(P) \neq \emptyset$ for every triangle T of G.*

$(a2)$ *Let $\vec{Q}$ be a cycle-dipacking in B. Then G has a Λ-factor P such that $\vec{F}(P,G) = \vec{Q}$.*

Proof (uses **2.4**). Obiously, $(a1.1) \Rightarrow (a1.2) \Rightarrow (a1.3)$. So all we need prove $(a1)$. Let $\vec{F} = \vec{F}(P,Q)$ and $v \in V(\vec{F})$. Let $d_{in}(v)$ and $d_{out}(v)$ denote the number of edges coming in v and going out of v in $\vec{F}$, respectively. By **2.4** and the fact that each vertex has been replaced by a triangle, there are three possibilities for every $v \in V(F)$:

$(c0)$ $d_{in}(v) = d_{out}(v) = 0$,

$(c1)$ $d_{in}(v) = 1$ and $d_{out}(v) = 1$, and

$(c2)$ $d_{in}(v) = 0$ and $d_{out}(v) = 3$.

If F has no vertices satisfying $(c2)$, then we are done. So let u be a vertex in $\vec{F}$ satisfying $(c2)$. Let S be a maximal directed path starting at u and ending at x. Clearly, $d_{in}(x) = 1$. Then x satisfies $(c1)$, and so $d_{out}(x) = 1$. Therefore $\vec{F}$ has an edge xy going out of x. Since S is a maximal directed path starting at u, clearly $y \in V(S - \{u, x\})$.

Then $d_{in}(y) = 2$, a contradiction. Finally, $(a2)$ follows from the fact that $G = B\{K_4, x\}$ and that every 2-edge matching covers the vertices of every triangle in K_4. $\square$

Below we give a construction that provides a big variety of graph-compositions with the properties similar to that of a special graph-composition in **4.1**.

4.2 *Let B and each A^u, $u \in V(B)$, be a cubic 3-connected graphs and $G = B\{(A^u, a^u) : u \in V(B)\}$. Suppose that*

$(h0)$ $v(B) = 0 \bmod 6$ *and each* $v(A^u) = 2 \bmod 6$*, and so* $v(G) = 0 \bmod 6$*,*

$(h1)$ *each $A^u - (N(a^u) \cup a^u \cup x)$ has no Λ-factor for every vertex x in $A^u - (N(a^u) \cup a^u)$ adjacent to a vertex in $N(a^u)$, and*

$(h2)$ *each $A^u - \{a^u, z\}$ has a Λ-factor for every $a^u z \in E(A^u)$, and*

$(h3)$ *each $A^u - W$ has a Λ-factor for every 5-vertex path W in A^u centered at a^u.*

Then

$(a1)$ *if P is a Λ-factor of G, then $\alpha^{-1}\{E(P) \cap E'(G)\}$ is the edge set of a Λ-factor Q of B and $\vec{F}(P, G) = \vec{Q}$,*

$(a2)$ *if P is a Λ-factor of G and if B and each A^u have no 3-blockades, then $|K \cap E(P)| \in \{1, 2\}$ for every 3-blockade K in G, and*

$(a3)$ *if $\vec{Q}$ is a Λ-difactor of B, then there exists a Λ-factor P of G such that $\vec{F}(P, G) = \vec{Q}$.*

Proof (uses **2.4**). Obviously, $(a2)$ follows from $(a1)$. We prove $(a1)$. Let $E'(P) = E(P) \cap E'(G)$ and P^u be the union of the components (3-vertex paths) of P meeting $D(A_u, G)$. By $(h0)$, $v(A^u) = 2 \bmod 6$. Therefore P^u satisfies $(a1.1)$ or $(a1.2)$ or $(a1.3)$ in **2.4**. By $(h1)$, P^u does not satisfy $(a1.3)$ in **2.4**. Thus $\alpha^{-1}(E'(P))$ is the edge set of a Λ-factor P' in B and the two edges of the same 3-vertex path L in P' are directed to the center vertex of L. Finally, $(a3)$ follows from assumption $(h2)$ and $(h3)$. $\square$

It is also easy to prove the following claim similar to that above when the assumption $v(A^u) = 2 \bmod 6$ is replaced by $v(A^u) = 4 \bmod 6$.

4.3 *Let B and each A^u, $u \in V(B)$, be a cubic 3-connected graphs and $G = B\{(A^u, a^u) : u \in V(B)\}$. Suppose that*

$(h0)$ $v(B) = 0 \bmod 6$ *and each* $v(A^u) = 4 \bmod 6$*, and so* $v(G) = 0 \bmod 6$*,*

$(h1)$ *each $A^u - T^u$ has no Λ-factor, where T^u is a tree subgraph of A^u containing a^u and such that $T^u - a^u$ is a 3-edge matching,*

$(h2)$ *each $A^u - a_u$ has a Λ-factor, and*

$(h3)$ *each $A^u - \Pi$ has a Λ-factor, where Π is a 4-vertex path in A^u with a^u being an inner vertex of Π.*

Then

$(a1)$ *if P is a Λ-factor of G, then $\vec{F}(P, G)$ is a cycle-dipacking in B,*

($a2$) *if P is a Λ-factor of G, then $|D(A_u, G) \cap E(P)| \in \{0, 2\}$ for every $u \in V(B)$, and*

($a3$) *if $\vec{Q}$ is a cycle-dipacking in B, then G has a Λ-factor P of G such that $\vec{F}(P, G) = \vec{Q}$.*

The proof of ($a1$) and ($a2$) in **4.3** is similar to that in **4.1**. The proof of ($a3$) in **4.3** is similar to that in **4.2**.

In the next two claims we give constructions of bricks satisfying the assumptions ($h0$) - ($h3$) in **4.2** and **4.3**, respectively.

Let $Z = Y(A^1, a^1; A^2, a^2; A^3, a^3)$ (see Fig. 2), where A^3 is the graph having two vertices z, a^3 and three parallel edges with the end-vertices z, a^3, and so $A^3 - a^3 = z$. Let $K_i = D(A_i, Z)$, and so K_i is the blockade with three edges having exactly one end-vertex in A_i, $i \in \{1, 2\}$.

Using **2.4**, it is easy to prove the following two claims.

4.4 *Let (Z, z) be a pair defined above. Suppose that $v(A^i) = 0 \bmod 6$ for $i \in \{1, 2\}$. Then $v(Z) = 2 \bmod 6$ and*

($a1$) *$Z - (N(z, Z) \cup z \cup v)$ has no Λ-factor for every vertex v in $Z - (N(z, Z) \cup z)$* (see also **2.4** ($a1.3$) and Fig. 3),

($a2$) *if $zz' \in E(Z)$ and P is a Λ-factor of $Z - \{z, z'\}$, then $|K_i \cap E(P)| \in \{1, 2\}$ for $i \in \{1, 2\}$, and*

($a3$) *if W is a 5-vertex path in Z with z being the center vertex of W and P is a Λ-factor of $Z - W$, then $|K_i \cap E(P)| \in \{1, 2\}$ for $i \in \{1, 2\}$.*

4.5 *Let (Z, z) be a pair defined above. Suppose that $v(A^i) = 4 \bmod 6$ for $i \in \{1, 2\}$. Then $v(Z) = 4 \bmod 6$ and*

($a1$) *if T is a tree subgraph of Z containing z and such that $T - z$ is a 3-edge matching, then $Z - T$ has no Λ-factor* (see also **2.4** ($a2.3$) and Fig. 4),

($a2$) *if P is a Λ-factor of $Z - z$, then $|K_i \cap E(P)| \in \{0, 2, 3\}$ for $i \in \{1, 2\}$, and if in addition, both A^1 and A^2 are isomorphic to K_4, then $|K_i \cap E(P)| = 2$ for $i \in \{1, 2\}$,*

($a3$) *if Π is a 4-vertex path in Z with z being an inner vertex of Π and P is a Λ-factor of $Z - \Pi$, then $|K_i \cap E(P)| \in \{0, 2\}$ for $i \in \{1, 2\}$, and*

($a4$) *if P is a Λ-factor of $Z - (N(z) \cup z)$, then $|K_i \cap E(P)| = 0$ for $i \in \{1, 2\}$.*

If (Z, z) satisfies the assumptions in **4.4**, then by **4.4**, the graph-composition $B\{Z, z\}$ satisfies the assumptions of **4.2**, and therefore has the properties ($a1$), ($a2$), and ($a3$) in **4.2**.

Similarly, if (Z, z) satisfies the assumptions in **4.5**, then by **4.5**, the graph-composition $B\{Z, z\}$ satisfies the assumptions of **4.3**, and therefore has the properties ($a1$), ($a2$), and ($a3$) in **4.3**.

Notice that if any claim in **3.1** is true, then there are infinitely many pairs (Z, z) satisfying the assumptions and, accordingly, the conclusions of **4.4**, as well as of **4.5**.

From **4.2** and **4.3** we have, in particular:

4.6 *Let* $\{i,j\} \in \{\{0,2\},\{1,2\}\}$. *Then there are infinitely many cubic 3-connected graphs* G *such that* $v(G) = 0 \bmod 6$ *and* $|E(P) \cap K| \in \{i,j\}$ *for every* Λ*-factor* P *of* G *and every 3-edge cut* K *of* G.

5 Λ-factor homomorphisms for graph-compositions

Let, as in Section 2, $G = B\{(A^u, a^u) : u \in V(B)\}$, $A_u = A^u - a^u$, and $E'(G) = E(G) \setminus \cup\{E(A_v) : v \in V(B)\}$. Let $N(a^u, A^u) = N^u$. As we mentioned in Section 2, there is a unique bijection $\alpha : E(B) \to E'(G)$ such that if $uv \in E(B)$, then $\alpha(uv)$ is an edge in G having one end-vertex in A_u and the other end-vertex in A_v.

Suppose that P is a Λ-factor of B. Let $V_s(P)$ be the set of vertices of degree s in P (and so $s \in \{1,2\}$). For a 3-vertex path uvw in P, let $A^u_1(P) = A^u - End(\alpha(uv))$ and let $A^v_2(P)$ be the graph obtained from A_v by adding two edges $\alpha(uv)$ and $\alpha(vw)$ together with their end-vertices.

Let $\Gamma(H)$ denote the set of Λ-factors of a graph H and $\Gamma(G,P)$ denote the set of Λ-factors Q of G such that $\alpha^{-1}(E'(G) \cap E(Q)\}) = E(P)$. Let $X \bigotimes Y$ denote the Cartesian product of sets X and Y (we assume that if $Y = \emptyset$, then $X \bigotimes Y = X$).

From **4.2** we have:

5.1 *Let* B *and each* A^u, $u \in V(B)$, *be cubic 3-connected graphs. Suppose that each* $v(A^u) = 2 \bmod 6$ *and each* (A^u, a^u) *satisfies the following conditions:*

$(h1)$ $A^u - (N^u \cup a^u \cup y)$ *has no* Λ*-factor for every vertex* y *in* $A^u - (N^u \cup a^u)$ *adjacent to a vertex in* N^u *and*

$(h2)$ $A^u - \{a^u, z\}$ *has a* Λ*-factor for every* $a^u z \in E(A(u))$, *and*

$(h3)$ $A^u - W$ *has a* Λ*-factor for every 5-vertex path* W *in* $A(u)$ *centered at* a^u.

Then :

$(\gamma 1)$ $\Gamma(G,P) \cap \Gamma(G,P') = \emptyset$ *for* $P, P' \in \Gamma(B)$, *and* $P \ne P'$,

$(\gamma 2)$ $\Gamma(G,P) = [\bigotimes\{\Gamma(A^v_1(P)) : v \in V_1(P)\}] \bigotimes [\bigotimes\{\Gamma(A^u_2(P)) : u \in V_2(P)\}]$, *and*

$(\gamma 3)$ $\Gamma(G) = \bigcup\{\Gamma(G,P) : P \in \Gamma(B)\}$,

and so G *has a* Λ*-factor if and only if* B *has a* Λ*-factor.*

Suppose that $\vec{C}$ is a cycle-dipacking in B. If uvw is a directed 3-vertex path in $\vec{C}$, then let $A^v_*(\vec{C})$ be the graph obtained from A_v by adding two edges $\alpha(uv)$ and $\alpha(vw)$ together with their end-vertices, a new vertex z, and a new edge wz.

Let $\Gamma(G, \vec{C})$ denote the set of Λ-factors Q in G such that $\vec{F}(G,Q) = \vec{C}$. Let $\mathcal{C}(B)$ denote the set of cycle-dipackings in B.

From **4.3** we have:

5.2 *Let B and each A^u, $u \in V(B)$, be cubic 3-connected graphs. Suppose that each $v(A(^u) = 2 \bmod 6$ and each (A^u, a^u) satisfies the following conditions:*

$(h0)$ each $v(A^u) = 4 \bmod 6$, and so $v(B) = 0 \bmod 6$ and $v(G) = 0 \bmod 6$,

$(h1)$ each $A^u - T^u$ has no Λ-factor, where T^u is a tree subgraph of A_u containing a^u and such that $T^u - a^u$ is a 3-edge matching,

$(h2)$ each $A^u - a_u$ has a Λ-factor, and

$(h3)$ each $A^u - \Pi$ has a Λ-factor, where Π is a 4-vertex path in A^u with a^u being an inner vertex of Π.

Then

$(\gamma 1)$ $\Gamma(G, \vec{C}) \cap \Gamma(G, \vec{C}') = \emptyset$ for $\vec{C}, \vec{C}' \in \mathcal{C}(B)$, $\vec{C} \neq \vec{C}'$,

$(\gamma 2)$ $\Gamma(G, \vec{C}) = [\bigotimes\{\Gamma(A^v_*(\vec{C}) : v \in V(\vec{C})\}] \bigotimes [\bigotimes\{\Gamma(A^u - a^u) : u \in V(B) - V(\vec{C})\}]$, and

$(\gamma 3)$ $\Gamma(G) = \bigcup\{\Gamma(G, \vec{C}) : \vec{C} \in \mathcal{C}(B)\}$.

As it is shown in Section 4, there are infinitely many pairs (A, a) satisfying the assumptions in **5.1**, as well as in **5.2**.

6 Cyclically 6-connected cubic graphs with special Λ-factor properties

In this section we describe a sequence (mentioned in Section 3, remark $(r2)$) of cyclically 6-connected graphs G with two disjoint 3-vertex paths L, L' such that $v(G) = 0 \bmod 6$ and $G - (L \cup L')$ has no Λ-factor.

We call a path T in a graph G a *thread* if the degree of each inner vertex of T in G is equal to two and the degree of each end-vertex of T in G is not equal to two.

Let C_s be a cycle with $9s$ vertices, $s \geq 1$, and let $\{L_k : k \in \{1, \ldots, 3s\}\}$ be a Λ-factor of C_s, where $L_i = (z^1_i z^2_i z^3_i)$. Let R_s be the graph obtained from C_s by adding the set $\{x^j_i : i \in \{1, \ldots, s\}, j \in \{1, 2, 3\}\}$ of $3s$ new vertices and the set $\{x^j_i z^j_i, x^j_i z^j_{i+s}, x^j_i z^j_{i+2s} : i \in \{1, \ldots, s\}, j \in \{1, 2, 3\}\}$ of $9s$ new edges (see Fig. 9 for an illustration of (R_1, L, L')). Let $X_i = \{x^1_i, x^2_i, x^3_i\}$.

Let $Q^i_{s-1} = R_s - X_i$. Then the vertices of 3-vertex paths L_i, L_{i+s}, and L_{i+2s} are of degree two in Q^i_{s-1}. Let T_i, T_{i+s}, and T_{i+2s} be the threads in Q^i_{s-1} containing L_i, L_{i+s}, and L_{i+2s}, respectively, and so each of these threads has exactly five vertices. Let R^i_{s-1} be the graph obtained from Q^i_{s-1} by replacing threads T_i, T_{i+s}, and T_{i+2s} by edges t_i, t_{i+s}, and t_{i+2s}. Then obviously, R^i_{s-1} is isomorphic to R_{s-1}.

Let Y_i be the subgraph of R_s induced by the vertex subset $V(L_i \cup L_{i+s} \cup L_{i+2s}) \cup X_i$. Then $R_s = Q^i_{s-1} \cup Y_i$, $C_s \cap Y_i = L_i \cup L_{i+s} \cup L_{i+2s}$, and the set K^i of edges having one vertex in Y_i and the other vertex in $R_s - Y_i$ is a 6-blockade in R_s, $s \geq 2$.

6.1 *Let R_s, $s \geq 1$, be the graph defined above, and so $v(R_s) = 12s$. Then*



Figure 9: (R_1, L, L'), $v(R_1) = 12$

(a1) *$R_s - (L \cup L')$ has no Λ-factor for every pair $\{L, L'\} \subset \{L_i, L_{i+s}, L_{i+2s}\}$, $i \in \{1, \dots, s\}$,*

(a2) *$c(R_s) = 6$ for $s \geq 2$ and $c(R_1) = 5$ (and for every 5-blockade M of R_1 one of the two components of $R_1 - M$ is a 5-cycle), and*

(a3) *$R_s - L - e$ has a Λ-factor for every edge e and every 3-vertex path L in $R_s - e$.*

Proof. The proof will consist of three parts.

(**p1**) We prove (a1). By symmetry, we can assume that $L = L_{1+s}$ and $L' = L_{1+2s}$. Let $G = R_s - (L \cup L')$. Suppose, on the contrary, that $G = R_s - (L \cup L')$ has a Λ-factor P. Then each edge $x_1^j z_1^j$, $j \in \{1, 2, 3\}$, is a dangling edge in G. Then $x_1^2 z_1^2$ belongs to a 3-vertex path L''. By symmetry, we can assume that $L'' = z_1^3 z_1^2 x_1^2$. Then x_1^1 is an isolated vertex in $G - L''$ (see, for example, Fig. 9, where $x^i = x_1^i$). Therefore P is not a Λ-factor of G, a contradiction.

(**p2**) We prove (a2) by induction on s. It is easy to check that R_1 has the properties described in (a2). So let $s \geq 2$. Since R_s has 6-cycles, $c(R_s) \leq 6$. So our goal is to prove that $c(R_s) = 6$. Let M be a minimum blockade in R_s. Obviously, each R_s is a connected graph, and so $M \neq \emptyset$.

Let $Q_{s-1} = Q_{s-1}^s$, and so the vertices of 3-vertex paths $L^1 = L_s$, $L^2 = L_{2s}$, and $L^3 = L_{3s}$ are of degree two in Q_{s-1}. Let $T^1 = T_s$, $T^2 = T_{2s}$, and $T^3 = T_{3s}$ be the threads in Q_{s-1} containing L^1, L^2, and L^3, respectively. Then graph R_{s-1} can be obtained from Q_{s-1} by replacing each thread T^j by a new edge t^j, $j \in \{1, 2, 3\}$. Let $Y = Y_s$. Then

$R_s = Q_{s-1} \cup Y$ and $C_s \cap Y = L^1 \cup L^2 \cup L^3$ and the edge set $K = K^s$ is a 6-blockade.

We know that $c(R_1) = 5$ and by the induction hypothesis, $c(R_{s-1}) = 6$ if $s \geq 3$.

Suppose first that M has an edge ab in $R_s - Y$. Then $ab \in E(R_{s-1})$, and so M contains a subset M' which is a blockade in Q_{s-1} separating a and b. If $|M'| = 6$, then $6 \leq |M'| \leq |M|$, and we are done. Since $ab \in E(R_{s-1})$, the sizes of minimum blockades separating a and b in Q_{s-1} and R_{s-1} are the same. So if $s \geq 3$, then by the induction hypothesis, $|M'| = 6$, and we are done.

So let $s = 2$ and $|M'| < 6$. Since $c(R_1) = 5$, $|M'|$ is equal to the size of a minimum blockade in R_1, and so $|M'| = 5$. Since no minimum blockade of R_1 is a cut in R_2, we have $5 = |M'| < |M|$, and so $|M| = 6$.

Now suppose that M has no edge in $R_s - Y$, i.e., $M \subseteq E(Y) \cup K$. Then it is easy to check (using the symmetries of Y) that every such blockade has six edges, and so $|M| = 6$.

(**p3**) Finally, we prove $(a3)$. Let, as in (**p2**), $Y = Y_s$ and $X = X_s$. Obviously, each Y_i is isomorphic to Y. It is easy to prove the following claims.

Claim 1. *$Y - L - e$ has a Λ-factor for every edge e in Y and for every 3-vertex path L in $Y - e$.*

Claim 2. *Let a and b be distinct vertices of degree two in Y that belong to different components of $Y - X$ and let $Y(ab)$ be the graph obtained from Y by adding a new edge ab. Let S be a 3-vertex path in $Y(ab)$ containing edge ab. Then $Y(ab) - S - e$ has a Λ-factor for every edge e in $Y(ab) - E(S)$.*

Let $D = E(C_s) - E(\cup\{L_k : k \in \{1, \dots, 3s\})$. Obviously, D is a matching and $R_s = \cup\{Y_i : i \in \{1, \dots, s\}\} \cup D$. Now, if P_i is a Λ-factor of Y_i, then clearly $\cup\{P_i : i \in \{1, \dots, s\}\}$ is a Λ-factor of R_s. Therefore from Claim 1 we have:

Claim 3. *$R_s - L - e$ has a Λ-factor for every edge e in R_s and every 3-vertex path L in $R_s - D - e$.*

Thus, it remains to consider a pair (F, e), where F is a 3-vertex path containing an edge in D and $e \in E(R_s - F)$. Since D is a matching, F has exactly one edge d in D. By symmetry of R_s, we can assume that $d = z_1^3 z_2^1$ and $F = z_1^3 z_2^1 v$, where z_1^3 and z_2^1 are vertices of degree two in Y_1 and Y_2, respectively, and $v \in \{x_2^1, z_2^2\}$.

For $z \in V(C_s)$, let $\alpha(z) = x$ if $xz \in E(R_s)$ and $x \in X$. Obviously, α is a function from $V(C_s)$ onto X.

(**p3.1**) Suppose first that $v = x_2^1$. Let $F_{2i-1} = z_{2i-1}^3 z_{2i}^1 \alpha(z_{2i}^1)$, and so $F_1 = F$. Let $M = \cup\{F_{2i-1} : \{i \in \{1, \dots, s\}\}$. By symmetry, we can assume that $e \notin E(M)$. Obviously, M is a Λ-packing in R_s and $Y_i \cap M$ consists of three vertices a_i, b_i, and x_i and one edge $a_i x_i$, where a_i and b_i have degree two in Y_i and belong to different components of $Y_i - X_i$ and $x_i \in X_i$. Let $S_i = x_i a_i b_i$. Then S_i is a 3-vertex path in $Y_i(a_i b_i)$. By Claim 2, $Y_i(a_i b_i) - S_i - e_i$ has a Λ-factor N_i for every edge e_i in $Y_i(a_i b_i) - E(S_i)$. Let $N = \cup\{N_i : i \in \{1, \dots, s\}\}$. Since $V(Y_i \cap M) = V(S_i)$, we have: $M \cup N$ is a Λ-factor of

R_s containing F and avoiding e.

(**p3.2**) Finally, suppose that $v = z_2^2$. Let $F_i = z_{2i-1}^3 z_{2i}^1 z_{2i}^2$, and so $F = F_1$. Let $M = \cup\{F_i : i \in \{1, \ldots, s\}\}$. By symmetry, we can assume that $e \not\in E(M)$. As in (**p3.1**), M is a Λ-packing in R_s and $Y_i \cap M$ consists of three vertices a_i, b_i, and x_i and one edge $a_i x_i$, where a_i and b_i have degree two in Y_i and belong to different components of $Y_i - X_i$ and $x_i \in X_i$. Therefore we are done by the same arguments as in (**p3.1**). $\square$

Acknowledgment. I am grateful to the referees for a very careful reading of the paper and very useful remarks. Also, many thanks to one of the referees for drawing my attention to recent papers [14–16].

References

[1] J.A. Bondy and U.S.R. Murty, *Graph Theory*, Springer, 2007.

[2] R. Deistel, *Graph Theory*, Springer-Verlag, New York, 2005.

[3] P. Hell and D. Kirkpatrick, Packing by complete bipartite graphs, *SIAM J. Algebraic Discrete Math.* **7** (1986), 199-209.

[4] A. Kelmans, Constructions of cubic bipartite and 3-connnected graphs without hamiltonian cycles. In *Analiz Zadach Formirovaniya i Vybora Alternativ*, VNIISI, Moscow **10** (1986) 64-72 (see also *Amer. Math. Soc. Transl.* (2) Vol. **158** (1994) 127-140).

[5] A. Kelmans, Packing k-edge trees in graphs of restricted vertex degrees, *J. Graph Theory*, **55** (2007) 306–324 (see also *DIMACS Research Report 2000–44*, Rutgers University (2000)).

[6] A. Kelmans, Packing k-paths in a cubic graph is NP-hard for $k \geq 3$, manuscript, 2001; presented at the DIMACS conference “Graph Partition”, July 2000.

[7] A. Kelmans, Packing 3-vertex paths in 2-connected graphs, *ArXiv: math.CO/ 0712.4151v1*, 26 December, 2007 (see also *RUTCOR Research Report RRR 21–2005*, Rutgers University (2005)).

[8] A. Kelmans, On Λ-packings in 3-connected graphs, *RUTCOR Research Report 23–2005*, Rutgers University (2005).

[9] A. Kelmans, Quasi Λ-factors in cubic Hamiltonian graphs, manuscript.

[10] A. Kelmans, Counterexamples to the cubic graph domination conjecture, *ArXiv: math.CO/0607512v1*, 24 July 2006.

[11] A. Kelmans, Packing 3-vertex paths in claw-free graphs, *ArXiv: math.CO/0711.3871v1*, 25 November 2007.

[12] A. Kaneko, A. Kelmans, T. Nishimura, On packing 3-vertex paths in a graph, *J. Graph Theory* **36** (2001) 175-197.

[13] A.K. Kelmans and D. Mubayi, How many disjoint 2-edge paths must a cubic graph have? *Journal of Graph Theory*, **45** (2004) 57–79 (see also *DIMACS Research Report 2000–23*, Rutgers University (2000)).

[14] A. Kosowski, M. Malafiejski, P. Zylinski, Tighter bounds on the size of a P_3-matching in a cubic graph, *Graphs and Combinatorics*, 24(5), (2008), 461-468.

[15] A. Kosowski, P. Zylinski, Packing three-vertex paths in 2-connected cubic graphs, *Ars Combinatoria* 89: (2008) 95-113.

[16] A. Kosowski, M. Malafiejski, and P. Zylinski, Parallel processing subsystems with redundancy in a distributed environment, PPAM, LNCS, 3911, (2005) 1002-1009.

[17] A.V. Kostochka and B.V. Stodolsky, On domination in connected cubic graphs, *Discrete Mathematics* 304 (2005) 749-762.

[18] B. Reed, Paths, stars, and the number three, *Combin. Probab. Comput.* 5 (1996) 277-295.

[19] D. West, *Introduction to Graph Theory*, Prentice Hall, 2001.